

# Early universe phase transitions within holographic composite Higgs model

Andrey Shavrin

under the supervision of Prof. Oleg Novikov  
Saint-Petersburg State University, Th. Phys. Dep.

Moscow International School on Particle Physics  
HSE Study Center “Voronovo”, Moscow

Mar. 4, 2024

# AdS/CFT correspondence

## Perturbation theory

$$\langle \phi \dots \phi \rangle = \langle \phi_0 \dots \phi_0 \rangle + \lambda \langle \phi_1 \dots \phi_1 \rangle + \lambda^2 \langle \phi_2 \dots \phi_2 \rangle + \dots + \text{non-analytical?} + \text{solitons?}$$

$$\text{QFT} \xrightarrow{\text{renormalization}} \text{CFT}$$

---

## AdS/CFT correspondence

$$\text{CFT} \xleftarrow{\text{AdS/CFT}} \text{dual theory}$$

$$\mathcal{Z}_{\text{CFT}}[J] \xleftrightarrow{\text{AdS/CFT}} \mathcal{Z}_{\text{AdS}} \xrightarrow{\text{quasiclassical}} e^{-S_{\text{AdS}}} \Big|_{\partial \text{AdS}}$$

correspondence      approximation

$$\text{strong coupled } \lambda_{\text{CFT}} \gg 1 \text{ with } \lambda_{\text{CFT}} \sim \frac{1}{\lambda_{\text{AdS}}}$$
$$\text{dual theory } \lambda_{\text{AdS}} \ll 1 \text{ weak coupled}$$

# Correlators within AdS/CFT

$$\text{Solutions of EoM: } \frac{\delta S_{\text{AdS}}[\psi]}{\delta \psi} = 0$$

$$\text{Near the conformal boundary: } \psi(x, z) \xrightarrow{z \rightarrow 0} z^{d-\Delta} \psi_0(x) + z^\Delta \psi_1(x)$$

$$\text{CFT sources: } J \sim \psi_0(x) \text{ with weight } d - \Delta$$

---

$$\text{Solutions: } \Rightarrow \mathcal{Z}_{\text{AdS}}|_{\partial\text{AdS}}[\psi_0] = \mathcal{Z}_{\text{CFT}}[J] = \int \mathcal{D}[\text{*fields of CFT*}] e^{-S_{\text{CFT}} - \mathcal{O} \cdot J}$$

$$G_n = \langle \mathcal{O} \dots \mathcal{O} \rangle = \left( \frac{\delta}{\delta J} \right)^n \log \mathcal{Z}[J] \Big|_{J=0} = - \left( \frac{\delta}{\delta \psi_0} \right)^n S_{\text{AdS}}[\psi] = \underbrace{- \left( \frac{\delta^n S_{\text{AdS}}^{\text{bulk}}}{\delta \psi_0^n} \right)}_{=0 \text{ due to EoM}} - \left( \frac{\delta^n S_{\text{AdS}}^{\text{border}}}{\delta \psi_0^n} \right)$$

# The price

AdS/CFT correspondence is a hypothesis in general sense.

Exact dualities: scalar theory / same in AdS<sub>5</sub> |  $\mathcal{N} = 4$  SYM / type IIB AdS<sub>5</sub> × S<sup>5</sup>

---

wanted CFT ⇒ Formulated FT in AdS (low energy limit of ST)

$$\mathcal{Z}_{\text{AdS}}|_{\partial\text{AdS}} = \int \mathcal{D}[???] e^{???) e^{-\mathcal{O}\cdot J}$$

Fields of CFT are unknown, action is unknown,  
but we know something (symmetries) about the sources  $J$  and the operators  $\mathcal{O}$

$$\mathcal{O} \in \mathbb{C}[\phi, \partial\phi, \partial^2\phi, \dots]$$

# Electroweak baryogenesis (Motivation)


Baryon asymmetry problem – matter more than anti-matter

Sakharov's conditions  $\Leftrightarrow$  first order phase transition (i.e. CPT-violation)

---


unbroken symmetry

$$\langle \phi \rangle = 0$$



broken symmetry

$$\langle \phi \rangle \neq 0$$



---

$SU(2)_L \times U(1)_Y \rightarrow U(1)_{em}$  is a crossover (not a PT); BSM physics?

# Composite Higgs model

$\mathcal{L} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{CH}} + \mathcal{L}_{\text{Int.}}$ ,  $\mathcal{L}_{\text{CH}}$  - strongly coupled with  $\mathcal{G}$  inner symmetry

$(\mathcal{G} \text{ invariant vacuum}) \xrightarrow{\text{spontaneous breaking}} (\mathcal{H} \text{ invariane vacuum}) \Rightarrow \text{Goldstone bosons} \ni \text{Higgs boson phase transition}$

---

$$\Sigma_{IJ} = \langle \bar{\Psi}_I \Psi_J \rangle = \xi^T \left[ \begin{pmatrix} 0_{4 \times 4} & 0 \\ 0 & \varsigma \end{pmatrix} + \eta_i \tilde{T}_i \right] \xi \xrightarrow[\text{low energy}]{\text{SO}(5) \rightarrow \text{SO}(4)} \begin{pmatrix} 0_{4 \times 4} & 0 \\ 0 & \varsigma \end{pmatrix} \Rightarrow \text{symmetry breaking}$$

$\Sigma_{IJ}$  is a condensate of the  $SO(5)$ -inn.sym. fundamental fields  $\Psi$ ;  
 $\xi$  is NB-bosons,  $\eta$  is “radial” fluctuations,  $\varsigma$  is background field

# Holographic Model

$$\text{Dual theory: } S_{\text{AdS}} = S_{\text{grav.}+\phi} + S_X + S_{\text{gauge}} + S_{\text{“SM”}} + S_{\text{int. “SM”}+\chi} \underbrace{\hspace{2cm}}_{\text{AdS/CFT}} S_{\text{SM}+\text{CH}}$$

$S_{\text{“SM”}}$  and  $S_{\text{int. “SM”}+\chi}$  shouldn't have a PT


---

$$\text{Matter sector: } S_X = \frac{1}{k_s} \int d^5x \sqrt{|g|} e^\phi \left[ \frac{1}{2} g^{ab} \text{Tr} \left( \nabla_a X^T \nabla_b X \right) - V_X(X) \right]$$

$$V_X(X) = \text{Tr} \left( -\frac{3}{2L^2} X^T X - \frac{\alpha}{4} (X^T X)^2 + L^2 \frac{\beta}{6} (X^T X)^3 + \mathcal{O}(X^8) \right)$$

$$X_{IJ} \propto \frac{\sqrt{N}}{2\pi} J_{IJ} z + \frac{2\pi}{\sqrt{N}} \Sigma_{IJ} z^3 + \dots, \quad J_{IJ} \text{ are sources for CH condensate } \Sigma_{IJ}$$

# Phase transition



$$\left. \frac{\partial V_{\text{eff}}^{\text{CH}}}{\partial \langle\varphi\rangle} \right|_{\langle\varphi\rangle_0} = 0 \Rightarrow \left( \langle\varphi\rangle_0, V_{\text{eff}} \big|_{\langle\varphi\rangle_0} \right)$$

Effective potential extremal values and the positions allow one to judge about PT:

- ❖ trivial minimum (vacuum) only  $\Rightarrow$  there is no PT;
- ❖ **non-trivial true vacuum with the potential barrier  $\Rightarrow$  1-st PT;**
- ❖ non-trivial true vacuum without a potential barrier  $\Rightarrow$  there is no PT.


The extrema of the effective quantum potential  $V_{\text{eff}}$ :  $T$  is the plasma temperature,  $\langle\varphi\rangle$  is the vacuum expectation.

*Unscaled schematic illustration! Data in real scale are at the backup slides.*



# Bubble free energy

Free energy of a bubble:  $F[V_{\text{eff}}]$  thin walls approximation  $4\pi R^2 \mu - \frac{3\pi}{4} R^3 (\mathcal{F}_{\text{out}} - \mathcal{F}_{\text{in}})$



$F_C \stackrel{\text{def}}{=} F(R_C)$ : if  $R > R_C$ , bubbles grow and **PT occurs**.

$\frac{\beta}{H_*} \sim \frac{F_C}{T} + \mathcal{O}(T)$   
 $1/\beta \sim \text{appear} \rightarrow \text{collide time}$   
 $1/H_* \sim \text{universe expansion}$


$v_4 \sim 10^{-1} \ll 1, \quad C \sim 1$

$$10^3 \gtrsim \frac{\beta}{H_*} \gtrsim 10^5$$

$$0.28 \gtrsim \frac{T(\gamma)}{m} < \frac{T(\gamma_{\min})}{m}$$

$$T \sim \frac{1}{\gamma} < \frac{1}{\gamma_{\min}} - \text{suggestion, } \gamma_{\min} \ll 1$$

# Observations



Picks of the GW spectrum estimated within Holographic Composed Higgs model. There is only the scalar part produced during initial collisions of the bubble walls (i.e. sound and turbulence contributions are not currently included in the rough estimate.)

# Large D limit

Dual theory:  $S_{\text{AdS}} = S_{\text{grav.}+\phi} + S_X + S_{\text{gauge}} + S_{\text{“SM”}} + S_{\text{int. “SM”}+X}$

$S_X$  is considered (without fluctuations);  $S_{\text{grav.}+\phi}$  and  $S_{\text{gauge}}$  are left

$$S_{\text{grav.}+\phi} = \frac{1}{l_p^3} \int d^5x \sqrt{|g|} e^{2\phi} \left[ -R + 2|\Lambda| - 4g^{ab} \partial_a \phi \partial_b \phi - V_\phi(\phi) \right]$$

---

$S_{\text{grav.}+\phi}$  within large D limit:  $g_{ab} = g_{ab}^{(0)} + \frac{1}{D} g_{ab}^{(1)} + \frac{1}{D^2} g_{ab}^{(2)} + \dots$

near-AdS<sub>d</sub> (Poincaré patch):  $A(z) = 1 + o(e^D)$ ,  $B(z) = 1 + o(e^D)$ ,  $f(z) = 1 - \left(\frac{z}{z_H}\right)^{D-1} + \mathcal{O}(e^D)$

$$ds^2 = A(z) \left( -f(z) dt^2 + \frac{dz^2}{f(z)} + B(z) d\Omega_{D-2} \right) \xrightarrow{D \rightarrow \infty} \frac{l^2}{D^2} (-y^2 d\tau^2 + dy^2) + o(e^{-D}) d\Omega_{D-2}$$

The simplest case for dilaton in AdS  $\phi'' - (\phi')^2 + m^2(\phi^2 + 2\phi) = 0$  no Lie symmetries

# Self-promotion

Thank you for the attention!



HELP DORA FIND THE SOURCE

# Effective field theory

$$Z[J] = \int \mathcal{D}\phi e^{-S-J\cdot\phi} =: e^{W[J]}$$

$$\Gamma[\langle\phi\rangle] = W[J] - \frac{\delta W[J]}{\delta J} \cdot J = \int_X d^d x \left( \underbrace{K_{\text{eff}}[\partial\langle\phi\rangle]}_{=0 \text{ if } \langle\phi\rangle=\text{const}} + V_{\text{eff}}[\langle\phi\rangle] \right) \quad \text{- effective action}$$

---

$$\text{Effective potential: } V_{\text{eff}} = \frac{1}{\text{Vol}_4} \Gamma$$

$$\text{Equation of motion (EoM): } \frac{\delta\Gamma}{\delta\langle\phi\rangle} = J \stackrel{\langle\phi\rangle=\text{const}}{=} \frac{\delta V_{\text{eff}}}{\delta\langle\phi\rangle} \stackrel{J=0}{=} 0 \quad \text{gives extrema condition}$$

# Effective potential

Extrema condition:  $V_{\text{eff}}|_{\text{extrema}} = V_{\text{eff}}|_{J=0} \Leftrightarrow G_0$ ;  
AdS/CFT:  $G_0 \Leftarrow$  boundary term of dual theory  $S_{\partial\text{AdS}}$

$$\text{Vol}_X V_{\text{eff}}|_{\text{extrema}} = G_0 = W[J=0] \stackrel{\text{AdS/CFT}}{=} S_{\text{AdS}}|_{\psi_0=0}$$

---

Extrema condition & duality:  $J = \psi_0 = 0$ ; duality:  $\langle \phi \rangle = \psi_1$

$$\frac{\delta V_{\text{eff}}}{\delta \langle \phi \rangle} \stackrel{\text{AdS/CFT}}{=} \frac{\delta}{\delta \psi_1} (S[\psi]|_{\partial\text{AdS}}) \Big|_{\psi_0=0} = 0 \quad \left( \begin{array}{c} \text{with assumption} \\ \langle \phi \rangle = \text{const} \end{array} \right)$$

gives vacuum expectation values:  $\{ \langle \phi \rangle_{\text{min } 1}, \langle \phi \rangle_{\text{min } 2}, \dots \}$  — possible vacuums

---

Extrema positions and values  $\{ (\langle \phi \rangle_{\text{min } i}, V_{\text{eff}}[\langle \phi \rangle_{\text{min } i}]) \} \Rightarrow$  phase transitions

# Holographic effective potential

$$\mathcal{Z}_{\text{CH}}[J] = \int \mathcal{D}\varphi \exp\left(-S[\varphi] - \int d^4x \varphi(x) J(x)\right) \stackrel{\text{def}}{=} e^{-W[J]}$$

$$\langle \varphi \rangle = \left. \frac{\delta W[J]}{\delta J} \right|_{J=0}, \quad \Gamma[\langle \varphi \rangle] = W[J] - \int d^4x \frac{\delta W[J]}{\delta J(x)} J(x) - \text{Effective Action}$$

$$\boxed{\text{EoM: } \frac{\delta \Gamma}{\delta \langle \varphi \rangle} = J}$$

extrema condition

Homogeneous Solution  $\Rightarrow \langle \varphi \rangle = \text{const}_{\mathbb{R}^{1,3}} \Rightarrow \Gamma = -\text{Vol}_4 V_{\text{eff}} - \text{Effective Potential}$

$\mathcal{Z}[J] \xrightarrow[\text{correspondence}]{\text{AdS/CFT}} \mathcal{Z}_{\text{AdS}} \xrightarrow[\text{approximation}]{\text{quasiclassical}} e^{-S_{\text{AdS}}}|_{\partial\text{AdS}} - \text{quasiclassical non-perturbative}$

$$\boxed{V_{\text{eff}} = -\frac{1}{\text{Vol}_4} S_{\text{AdS}}|_{\partial\text{AdS}}} - \text{boundary term of the bulk theory defines } \textit{quantum} \text{ effective potential}$$

# Holographic model

$\mathcal{L}_{\text{CH}}$  - strongly coupled  $\Rightarrow$  consider  $N \gg 1 \Rightarrow \mathcal{Z}_{\text{CH}}[J] = \mathcal{Z}_{\text{AdS}}[J]$

The dual theory:  $\mathcal{Z}_{\text{AdS}}[J] \sim \exp(-S_{\text{AdS}}[J])$  is weakly coupled  $\Rightarrow$  quasiclassical limit

---

**The asymptotic behavior** near the conformal border  $\partial\text{AdS}$  of the dual theory fields **defines the sources** of the CH operators (i.e. the **correlator functions**)

$$X_{IJ} \xrightarrow{z \rightarrow 0} \frac{\sqrt{N}}{2\pi} J_{IJ} z + \frac{2\pi}{\sqrt{N}} \Sigma_{IJ} z^3 + \dots \quad X_{IJ} : \text{AdS}_5 \xleftrightarrow{\text{dual}} \Sigma_{IJ} : \mathbb{R}^{1,3}$$

---

Holography is the **duality** between **strongly coupled** theory on the border and **weakly coupled** (quasiclassical) bulk theory.

$$F = -T \log \mathcal{Z}_{\text{CH}} \sim T S_{\text{AdS}} \propto \text{Vol}_4 \cdot \mathcal{F} \quad \text{In homogeneous case } (\chi = \chi(z)): \mathcal{F} \propto V_{\text{eff}}[\chi]$$



# Action of the holographic model

$$S_{\text{tot}} = S_{\text{grav}+\phi} + S_X + S_A + S_{\text{SM}} + S_{\text{int}}, \quad S_A = -\frac{1}{g_5^2} \int d^5x \sqrt{|g|} e^\phi g^{ac} g^{bd} F_{ab} F_{cd}$$

$$S_{\text{grav}+\phi} = \frac{1}{l_p^3} \int d^5x \sqrt{|g|} e^{2\phi} \left[ -R + 2|\Lambda| - 4g^{ab} \partial_a \phi \partial_b \phi - V_\phi(\phi) \right], \quad a, b = 0, \dots, 4$$

$$S_{\text{int}} = \epsilon^4 \int_{z=\epsilon} d^4x \sqrt{|g^{(4)}|} \left[ c_Y B_\mu \text{Tr} (T_Y A^\mu) + c_W W_{k,\mu} \text{Tr} (T_k A^\mu) + \mathcal{L}_\psi \right]$$

$$S_X = \frac{1}{k_s} \int d^5x \sqrt{|g|} e^\phi \left[ \frac{1}{2} g^{ab} \text{Tr} (\nabla_a X^T \nabla_b X) - V_X(X) \right], \quad \nabla_a X = \partial_a X + [A_a, X], \quad A_a = 0$$

$$V_X(X) = \text{Tr} \left( -\frac{3}{2L^2} X^T X - \frac{\alpha}{4} (X^T X)^2 + L^2 \frac{\beta}{6} (X^T X)^3 + \mathcal{O}(X^8) \right)$$

$$L \cdot X_{IJ} \sim \frac{\sqrt{N}}{2\pi} J_{IJ} \tilde{z} + \frac{2\pi}{\sqrt{N}} \Sigma_{IJ} \tilde{z}^3 + \dots$$

# Geometry

$$S_{\text{grav}+\phi} = \frac{1}{l_p^3} \int d^5x \sqrt{|g|} e^{2\phi} \left[ -R + 2|\Lambda| - 4g^{ab} \partial_a \phi \partial_b \phi - V_\phi(\phi) \right], \quad a, b = 0, \dots, 4$$

$$ds^2 = \frac{L^2}{\tilde{z}^2} A(\tilde{z})^2 \left( f(\tilde{z}) d\tau^2 + \frac{d\tilde{z}^2}{f(\tilde{z})} + d\vec{x}^2 \right), \quad \phi = \phi(\tilde{z})$$

$$f = 1 - \frac{\tilde{z}^4}{z_H^4}, \quad \phi = \tilde{\phi}_2 \tilde{z}^2, \quad z_H = \frac{1}{\pi T}.$$

# Temperature estimations

Experimental restrictions  $\Leftrightarrow$  mass of the **lightest predicted** particle.

$$\Sigma_{IJ} = \xi^\top \left[ \begin{pmatrix} 0_{4 \times 4} & 0 \\ 0 & \varsigma \end{pmatrix} + \eta_i \tilde{T}_i \right] \xi \xrightarrow[\text{dual to}]{\text{AdS/CFT}} \chi_{IJ} \rightarrow \begin{pmatrix} 0_{4 \times 4} & 0 \\ 0 & \chi \end{pmatrix}$$

$m_\eta \sim m_{\delta\chi}$  fluctuation mass  $\sim$  slope of the “hat”.

$$\chi(z) \rightarrow \chi(z) + \delta\chi(t, \vec{x}, z) \Rightarrow \text{EoM}_z[\chi] \rightarrow \text{EoM}_{t, \vec{x}, z}[\chi + \delta\chi] \xrightarrow{\partial \text{AdS}} \boxed{2m^2 = \phi_2}$$

---

$$T = \frac{1}{\pi} \frac{1}{z_H} \Rightarrow T = \frac{m}{\pi} \sqrt{\frac{2}{\phi_2}}, \quad z_H^2 = \frac{\phi_2}{2m^2}$$

# Gravitational Waves

The spectrum of the gravitational waves can be estimated as  
(within the approach of **relativistic** velocity of the bubble walls  $v_w \sim 1$ )

$$\Omega_{\text{GW}} h^2 = 1.67 \cdot 10^{-5} \kappa \Delta \left( \frac{\beta}{H_*} \right)^{-2} \left( \frac{\alpha}{1 + \alpha} \right)^2 \left( \frac{g_*}{100} \right)^{-\frac{1}{3}}$$

**Only scalar waves! Sound waves and turbulence are not included!**

We **estimate** only scalar waves produced during initial collisions.

$$f_0 = 1.65 \cdot 10^{-5} \text{Hz} \cdot \frac{f_*}{\beta} \frac{\beta}{H_*} \frac{T}{0.1 \text{TeV}} \left( \frac{g_*}{100} \right)^{\frac{1}{6}} \text{Hz}$$

$(\Omega_{\text{GW}} h^2, f_0)$ -curve is the estimation GW amplitude (peak value).

It does not contain the spectral shape  $S(f_0)$  (in this case  $S(f_0 = f_0^{\text{peak}}) = 1$ ).

# “Extrema” curves

$$\frac{\delta S_\chi}{\delta \chi} = 0 \Rightarrow \chi \xrightarrow{z \rightarrow 0} Jz + \left( \sigma - \left( \frac{3}{2} J^3 + \phi_2 J \right) \log z \right) z^3 + o(z^5) - \text{give the sources for CFT operators}$$

Knowing the *extrema* of the effective potential and its *values* at these points, we can judge about the phase transition

$$V_{\text{eff}} = -\frac{1}{\text{Vol}_4} S_\chi \Big|_{\partial \text{AdS}} \Rightarrow \text{from EoM for effective action: } \text{Vol}_4 \frac{\delta V_{\text{eff}}}{\delta \langle \varphi \rangle} = J \Rightarrow \text{extrema condition is absence of sources} \Rightarrow J = 0$$

$$\underbrace{\chi \xrightarrow{z \rightarrow 0} \sigma z^3 + o(z^5)}_{\text{“extreme” solutions}} \text{ must give } \underbrace{\frac{\delta V_{\text{eff}}}{\delta \langle \varphi \rangle} = 0}_{\text{extrema}} \Rightarrow \text{a new condition for } \phi_2 \text{ and } \langle \varphi \rangle$$

$$T \sim \frac{1}{\sqrt{\phi_2}}, \quad \frac{\delta V_{\text{eff}}}{\delta \langle \varphi \rangle} = 0 = \frac{\delta}{\delta \sigma} S_\chi [\chi_{\text{Sol.}}(z; J, \sigma)] \Big|_{J=0} \Rightarrow \{\sigma_1, \dots, \sigma_n\} - \text{extrema}$$

# Nucleation ratio

Baryogenesis generates enough asymmetry (enough efficient)  
if there is one bubble per Hubble volume

$$\underbrace{\text{Nucleation Ratio: } AT^4 e^{-\frac{F_C}{T}}}_{\text{Bubbles produced per time} \times \text{space volume}} \sim \underbrace{H^4(T) = \left(\frac{T^2}{M_{\text{Pl}}}\right)^4}_{1/(\text{Hubble time} \times \text{volume})} \quad \text{— Expansion of the Universe}$$

---

$F = F[\langle\varphi\rangle, R]$  – Free energy of the bubble;  $R$  is the radius of the bubble

Hubble horizon (time, volume, radius) — speed of receding object behind it is greater than the speed of light (Don't confuse with cosmological horizon)

Bubble appears with a certain size. It defines with “micro-physics”.


If its radius is greater, then critical one  $\left. \frac{\partial F}{\partial R} \right|_{R_C \stackrel{\text{def}}{=} R}$ , the bubble grows. Otherwise, it bursts.

It gives  $F_C \stackrel{\text{def}}{=} F(R_C)$  and defines nucleation ratio and “viability of the model”.

# Estimations of the nucleation ratio

$$\Gamma \sim H_*^4, \quad \frac{\Gamma}{m^4} \sim \frac{H_*^4}{m^4} \propto \frac{m^4}{M_{\text{Pl}}^4};$$

$F_C$  is defined with an error, so  $e^{\frac{F_C}{T}}$  has large error



# Potentials of CFT and the dual theory

$$V_\chi = a_2\chi^2 + a_4\chi^4 + a_6\chi^6, \quad a_2 < 0, a_4 < 0, a_6 > 0 \quad \text{no barrier}$$

$$V_{\text{eff}} = b_2\langle\varphi\rangle^2 + b_4\langle\varphi\rangle^4 + b_6\langle\varphi\rangle^6, \quad b_2 > 0, b_4 < 0, b_6 > 0 \quad \text{there's a barrier}$$

in details:

- $V_{\text{eff}} = V_{\text{eff}}[\langle\varphi\rangle]$  describes a quantum objects at the border.  $V_\chi$  is a dual classical potential in the bulk.
- $V_{\text{eff}} = -\frac{1}{\text{Vol}_4} S_{\text{AdS}} \Big|_{\partial\text{AdS}}$  includes the solutions of the EoM  $\frac{\delta S_\chi}{\delta\chi=0}$  in bulk. In other words,  $V_{\text{eff}}$  includes physics of AdS



# Conditions for the dual theory potential

$V_\chi(\chi) = \frac{m^2}{2}\chi^2 - \frac{D}{4L^2}\lambda\chi^4 + \frac{\lambda^2\gamma}{6L^2}\chi^6$  is the expansion of a more general theory

Suggestions:

- ❖ The potential  $V_\chi$  always has true vacuum with  $E_{\min} (V_\chi \xrightarrow{\chi \rightarrow \pm\infty} \infty)$ . So we may use any even power  $\chi^n$  instead of the last term  $\chi^6$ .
- ❖ The expansion of  $V_\chi$  has certain sign of the second term  $\lambda > 0$  (the first one  $m^2$  chosen for the theory to be conformal in AdS).
- ❖ Higher orders of the expansion don't give new minima at the considered temperatures.

---

The certain parametrization has been chosen with respect to the “symmetries”

“Scale invariance”,  
defining  
the coefficients :  $L \rightarrow L'$   
 $\chi \rightarrow \sqrt{\lambda}\chi$  ; Conformality near  
the AdS border :  $\Delta_- = 1$   
 $\Delta_+ = 3 \Rightarrow m^2 = -\frac{D}{3L^2}$   
 (“correct” conformal weights)

*D* is for the Large *D* limit. But its usage doesn't give any results.  
(to keep interaction constants finite at  $D \rightarrow \infty$ )

# SM - CH model interactions

$$F \stackrel{\text{thin walls}}{\text{approximation}} 4\pi R^2 \mu - \frac{3\pi}{4} R^3 (\mathcal{F}_{\text{out}} - \mathcal{F}_{\text{in}}) - \text{physical units are required}$$

- Fix the Parameters (Interaction with Standard Model – bulk gauge fields)
- Physical Units (Infrared Regularization and finite temperature – “radial” heavy fluctuations)

$$W_\mu^\alpha J_L^{\alpha\mu} + B_\mu J_Y^\mu \Leftrightarrow J^\mu \sim A^M - \text{bulk } \mathcal{G} \text{ gauge field}$$

---

The physical values can be estimated without gauge field:

$$\Sigma_{IJ} = \langle \bar{\Psi}_I \Psi_J \rangle = \xi^\top \left[ \begin{pmatrix} 0_{4 \times 4} & 0 \\ 0 & X \end{pmatrix} + \eta_i \tilde{T}_i \right] \xi \Leftrightarrow \frac{1}{T} \propto \sqrt{\phi_2} \sim \mu_{\text{IR}} \sim m_\eta \gtrsim 10 \text{ TeV}$$

$$m_\eta \Leftarrow X \rightarrow X + \delta X - \text{correction of the background field} \Rightarrow \eta - \text{pNG boson}$$

# CH gauge field

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{CH}} + B_\mu \text{Tr} \left( T_Y \hat{J}^\mu \right) + W_{k,\mu} \text{Tr} \left( T_k \hat{J}^\mu \right) + \sum_r \bar{\psi}_r \mathcal{O}_r + \text{h.c.}$$

$\underbrace{\hspace{15em}}_{=\mathcal{L}_{\text{interactions}}}$

$$SO(5) \times U(1) : A_M = A_M^K T^K + A_{M,Y} T_Y$$

$$SO(5) \rightarrow SO(4) : \underbrace{A_M^K T^K}_{\in SO(5)} \rightarrow \underbrace{A_M^a T^a}_{\in SO(4)} + \underbrace{A_M^i T^i}_{\in SO(5)/SO(4)}$$

$$SO(4) \cong SU(2) \times SU(2) : A_M^K T^K = \underbrace{A_M^{k,L} T_L^k}_{\in SU(2)_L} + \underbrace{A_M^{k,R} T_R^k}_{\in SU(2)_R}$$

conserved currents:  $\hat{J}_\mu \xleftrightarrow{\text{dual}} A_\mu(t, x, z) \Big|_{z=0}^{\partial \text{AdS}}$ ,    holographic gauge:  $A_z = 0$

$$\mathcal{O} \xleftrightarrow{\text{dual}} \mathcal{J}(A, \Psi, \phi, \dots) \text{ — composite operators of the CH fields}$$